

# Interior's Recommendations to the SWRCB re: DCC Gate Operations

- Interior will present information from past fishery studies that indicate diversion into the interior Delta lowers juvenile salmon survival. Closing the Delta Cross Channel would be an action that is intended to reduce the number diverted.
- DWR will explain how the salmon decision tree is used to protect juvenile salmon on a real-time basis.
- Interior recommends that the salmon decision tree and DAT CALFED Ops process continue to be used to provide fish protection related to the DCC gate operations
- When the DCC studies have been completed and DCC gate operations alternatives have been evaluated (estimated 5-7 years) Interior will address the SWRCB with specific recommendations re: the WQCP DCC gate operations.


# Salmon Conceptual Model and Basis for DCC Closures

SWRCB Periodic Review

November 15, 2004

Pat Brandes - USFWS

# Conceptual Model of juvenile salmon migration through the Delta




-Sac Basin salmon move into interior Delta through the open DCC and GS

- in the interior Delta their survival is lower  
- and a function of exports

- Vulnerable to entrainment during emigration

# Conceptual Model of juvenile salmon migration through the Sacramento Delta


-Sac Basin salmon move into interior Delta through the open DCC and GS



**Flood Tide**

**Gates open**


Dec. 13, 2001 at 03:20 PM PST



**EBB Tide**

**DCC gates closed**

# % Freeport flow diverted into interior Delta with DCC gates open and closed




# Conceptual Model of juvenile salmon migration through the Sacramento Delta

-Sac Basin salmon move into interior Delta through the open DCC and GS


-More move into interior Delta with gates open

**-Survival is lower in the interior Delta**



**Marked juvenile salmon releases made on Sac River above and below the DCC and GS**


# Survival indices to Chipps Island of marked juvenile salmon released upstream and downstream of the Delta Cross Channel and Georgiana Slough



**Upstream < downstream (p<0.05)**


Data would infer additional mortality of upstream group due to some entering DCC and GS

# Survival indices to Chipps Island of marked juvenile salmon released upstream and downstream of the Delta Cross Channel and Georgiana Slough




**Upstream < downstream (p<0.05)**

Data would infer additional mortality of upstream group due to some entering GS



**Release sites for marked salmon released on Sacramento River (Ryde and Isleton) and interior Delta (Georgiana Slough and Lower Mokelumne)**


# Survival indices to Chipps Island



Fall GS < Ryde (p < 0.05)

Late-fall GS < Ryde (p < 0.05)

# Recovery Rates in the ocean fishery



Fall GS < Ryde (p < 0.05)

Late-fall GS < Ryde (p < 0.05)

Ryde/Isleton


# Conceptual Model of juvenile salmon migration through the Sacramento Delta

-Sac Basin salmon move into interior Delta through the open DCC and GS


-More move into interior Delta with gates open

-Survival is lower In the interior Delta

-and a function of exports



Relationship between GS/Ryde survival ratio and CVP/SWP exports with 95% confidence intervals



Relationships between GS/Ryde survival ratio and CVP/SWP exports

# Conceptual Model of juvenile salmon migration through the Sacramento Delta




- Sac Basin salmon move into interior Delta through the open DCC and GS
- More move into interior Delta with gates open
- In the interior Delta, their survival is lower
- and a function of exports

**Statistical Modeling of fall run CWT recoveries is supportive of this conceptual model**



# OLDER JUVENILES

2001-2002












# NUMBER OF FRY/SMOLT CHINOOK RECOVERED IN THE SACRAMENTO RIVER AND DELTA

## 2000-2001



# **The basis of DCC fish protective actions in the Delta for juvenile salmon**

**is based on evidence that indicates:**

- survival in the Delta is lower in the interior Delta**
- and that with the DCC gates closed a lower percent of water and presumably juvenile salmon is diverted into the interior Delta**

**This action will likely increase the survival of juvenile salmon through the Delta**


**The timing of action is designed to increase survival during periods of high abundance of the various races/runs**

# Interior's Recommendations to the SWRCB re: DCC Gate Operations

- Interior presented information from past fishery studies that indicate diversion into the interior Delta lowers juvenile salmon survival. Closing the Delta Cross Channel would be an action that is intended to reduce the number diverted.
- DWR will explain how the salmon decision tree is used to protect juvenile salmon on a real-time basis.
- Interior recommends that the salmon decision tree and DAT CALFED Ops process continue to be used to provide fish protection related to the DCC gate operations
- When the DCC studies have been completed and DCC gate operations alternatives have been evaluated (estimated 5-7 years) Interior will address the SWRCB with specific recommendations re: the WQCP DCC gate operations.




Survival indices to Chipps Island of marked juvenile salmon released upstream and downstream of the Delta Cross Channel and Georgiana Slough with DCC gates open and closed




Nov. 15, 2001 at 11:43 AM PST

NATURAL RESOURCE SCIENTISTS, INC.  
**USGS**  
science for a changing world



GS/Ryde Survival Ratio (with 95% confidence intervals)



Exports in cfs (for 3 days after release)